

Naval Research Laboratory

Washington, DC 20375-5000

DTIC FILE COPY



2

NRL Report 9127

AD-A197 822

A Note on Incomplete Integrals of Cylindrical Functions

ALLEN R. MILLER

Engineering Services Division

June 21, 1988

DTIC
ELECTE
AUG 08 1988
S E D

Approved for public release; distribution unlimited.

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b RESTRICTIVE MARKINGS		
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b DECLASSIFICATION/DOWNGRADING SCHEDULE					
4 PERFORMING ORGANIZATION REPORT NUMBER(S) NRL Report 9127			5 MONITORING ORGANIZATION REPORT NUMBER(S)		
6a NAME OF PERFORMING ORGANIZATION Naval Research Laboratory		6b OFFICE SYMBOL (If applicable) Code 2303	7a NAME OF MONITORING ORGANIZATION		
6c ADDRESS (City, State, and ZIP Code) Washington, DC 20375-5000			7b ADDRESS (City, State, and ZIP Code)		
8a NAME OF FUNDING/SPONSORING ORGANIZATION		8b OFFICE SYMBOL (If applicable)	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c ADDRESS (City, State, and ZIP Code)			10 SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO	PROJECT NO	TASK NO
11 TITLE (Include Security Classification) A Note on Incomplete Integrals of Cylindrical Functions					
12 PERSONAL AUTHOR(S) Miller, A.R.					
13a TYPE OF REPORT Final		13b TIME COVERED FROM _____ TO _____		14 DATE OF REPORT (Year, Month, Day) 1988 June 21	
15 PAGE COUNT 11					
16 SUPPLEMENTARY NOTATION					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB GROUP			
			Hypergeometric functions		
			Cylindrical functions		
19 ABSTRACT (Continue on reverse if necessary and identify by block number) Representations for incomplete Lipschitz-Hankel integrals of cylindrical functions are given in closed form by using Kampé de Fériet double hypergeometric functions. In addition, reduction formulas for the Kampé de Fériet functions associated with these integrals are provided for some cases.					
20 DISTRIBUTION AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a NAME OF RESPONSIBLE INDIVIDUAL Allen R. Miller			22b TELEPHONE (Include Area Code) (202) 767-2215		22c OFFICE SYMBOL Code 2303

DD Form 1473, JUN 86

Previous editions are obsolete
S/N 0102-1E-014-6603

SECURITY CLASSIFICATION OF THIS PAGE

CONTENTS

INTRODUCTION	1
PRELIMINARY RESULTS AND DEFINITIONS	1
REPRESENTATIONS FOR $C_{e_{\mu, \nu}}(a, z)$, $C_{s_{\mu, \nu}}(a, z)$, $C_{c_{\mu, \nu}}(a, z)$	3
REDUCTION FORMULAS FOR L AND Q	5
APPLICATIONS	5
SUMMARY	7
REFERENCES	7

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



A NOTE ON INCOMPLETE INTEGRALS OF CYLINDRICAL FUNCTIONS

INTRODUCTION

The class of cylindrical functions C includes Bessel functions of the first kind J , modified Bessel functions I , Bessel functions of the second kind or Neumann functions Y (or N), Bessel functions of imaginary argument or MacDonald functions K , and Bessel functions of the third kind that include Hankel functions of the first and second kind, $H^{(1)}$ and $H^{(2)}$.

The general incomplete Lipschitz-Hankel integral of cylindrical functions $C_\nu(z)$ is defined as the function of two complex variables:

$$C_{e_{\mu,\nu}}(a, z) \equiv \int_0^z e^{at} t^\mu C_\nu(t) dt. \quad (1)$$

Here the symbol e denotes the presence of the exponential function and μ, ν may be complex. Analogously, we define integrals that contain the functions $\sin(at)$ and $\cos(at)$ in place of $\exp(at)$:

$$C_{s_{\mu,\nu}}(a, z) \equiv \int_0^z \sin(at) t^\mu C_\nu(t) dt \quad (2)$$

$$C_{c_{\mu,\nu}}(a, z) \equiv \int_0^z \cos(at) t^\mu C_\nu(t) dt. \quad (3)$$

To assure convergence of $C_{e_{\mu,\nu}}(a, z)$ and $C_{c_{\mu,\nu}}(a, z)$, it is necessary that $\operatorname{Re}(\mu + 1) > |\operatorname{Re} \nu|$ when $C = K, Y, H^{(1)}, H^{(2)}$; $\operatorname{Re}(1 + \mu + \nu) > 0$ when $C = I, J$. When $\mu = \nu$, we define, for example, $C_{e_{\mu,\mu}} \equiv C_{e_\mu}$ where for convergence $\operatorname{Re} \mu > -1/2$ for all C .

Integrals of the type given by Eqs. (1) to (3) occur very often in applied mathematics. Agrest and Maksimov [1] have found representations for $C_{e_{\mu,\nu}}(a, z)$, $C_{s_{\mu,\nu}}(a, z)$, and $C_{c_{\mu,\nu}}(a, z)$ using incomplete cylindrical functions. In this report we give representations for $C_{e_{\mu,\nu}}(a, z)$, $C_{s_{\mu,\nu}}(a, z)$, and $C_{c_{\mu,\nu}}(a, z)$ using only the Kampé de Fériet double hypergeometric functions $F_{2:1:0}^{0:2:1}[x, y]$.

PRELIMINARY RESULTS AND DEFINITIONS

To begin, we summarize some results that are found in Ref. 2, p. 85: Let a and b be arbitrary constants,

$$\mathbf{F}_\nu(z) \equiv aI_\nu(z) + be^{i\nu\pi}K_\nu(z)$$

$$\mathbf{G}_\nu(z) \equiv aJ_\nu(z) + bY_\nu(z)$$

$$\alpha \equiv \begin{cases} i: \mathbf{H} = \mathbf{F} \\ 1: \mathbf{H} = \mathbf{G} \end{cases} \quad \beta \equiv \begin{cases} 1: \mathbf{H} = \mathbf{F} \\ 0: \mathbf{H} = \mathbf{G} \end{cases}$$

Then

$$\int_0^z t^\mu \mathbf{H}_\nu(t) dt = e^{-\frac{\pi}{2} i \beta \mu} [(\mu + \nu - 1) z \mathbf{H}_\nu(z) s_{\mu-1, \nu-1}(\alpha z) + (2\beta - 1) \alpha z \mathbf{H}_{\nu-1}(z) s_{\mu, \nu}(\alpha z)], \quad (4)$$

where the Lommel functions $s_{\mu, \nu}$ are given by

$$s_{\mu, \nu}(z) = \frac{z^{\mu+1}}{(\mu - \nu + 1)(\mu + \nu + 1)} {}_1F_2 \left[1; \frac{\mu - \nu + 3}{2}, \frac{\mu + \nu + 3}{2}; \frac{-z^2}{4} \right]. \quad (5)$$

Now defining

$$\xi \equiv \begin{cases} 1: & C = I, K \\ -1: & C = H, J, Y \end{cases} \quad \eta \equiv \begin{cases} 1: & C = K \\ -1: & C = H, I, J, Y \end{cases}$$

we may deduce from Eqs. (4) and (5) the result

$$\begin{aligned} \int_0^z t^\mu C_\nu(t) dt &= \frac{z^{\mu+1}}{\mu - \nu + 1} \left\{ C_\nu(z) {}_1F_2 \left[1; \frac{\mu - \nu + 3}{2}, \frac{\mu + \nu + 1}{2}; \frac{\xi z^2}{4} \right] \right. \\ &\quad \left. + \frac{\eta z C_{\nu-1}(z)}{\mu + \nu + 1} {}_1F_2 \left[1; \frac{\mu - \nu + 3}{2}, \frac{\mu + \nu + 3}{2}; \frac{\xi z^2}{4} \right] \right\}. \end{aligned} \quad (6)$$

We define the Kampé de Fériet double hypergeometric functions L and Q and give associated generating relations [3, 4]:

$$\begin{aligned} L[\alpha, \beta; \gamma, \delta; x, y] &\equiv F_{2,0,0}^{0,1,1} \left[\begin{matrix} \text{---}; & \alpha; & \beta; \\ \gamma, \delta; & \text{---}; & \text{---}; \end{matrix} \middle| x, y \right], \quad |x| < \infty, \quad |y| < \infty \\ Q[\alpha, \beta, \gamma; \mu, \nu, \lambda; x, y] &\equiv F_{2,1,0}^{0,2,1} \left[\begin{matrix} \text{---}; & \alpha, \beta; & \gamma; \\ \mu, \nu; & \lambda; & \text{---}; \end{matrix} \middle| x, y \right], \quad |x| < \infty, \quad |y| < \infty \\ L[\alpha, \beta; \gamma, \delta; x, y] &= \sum_{m=0}^{\infty} \frac{(\alpha)_m}{(\gamma)_m (\delta)_m} \frac{x^m}{m!} {}_1F_2[\beta; m + \gamma, m + \delta; y] \\ Q[\alpha, \beta, \gamma; \mu, \nu, \lambda; x, y] &= \sum_{m=0}^{\infty} \frac{(\alpha)_m (\beta)_m}{(\mu)_m (\nu)_m (\lambda)_m} \frac{x^m}{m!} {}_1F_2[\gamma; m + \mu, m + \nu; y]. \end{aligned} \quad (7)$$

It is easy to see that the function L is a special case of Q :

$$Q[\alpha, \lambda, \beta; \gamma, \delta, \lambda; x, y] = L[\alpha, \beta; \gamma, \delta; x, y].$$

For brevity we define the parameter lists

$$A_1(\mu, \nu) \equiv \frac{\mu + \nu + 1}{2}, \frac{\mu - \nu + 1}{2}, 1; \frac{\mu + \nu + 1}{2}, \frac{\mu - \nu + 3}{2}, \frac{1}{2}$$

$$A_2(\mu, \nu) \equiv \frac{\mu + \nu + 1}{2}, \frac{\mu - \nu + 1}{2}, 1; \frac{\mu + \nu + 3}{2}, \frac{\mu - \nu + 3}{2}, \frac{1}{2}$$

$$B_1(\mu, \nu) \equiv \frac{\mu + \nu + 2}{2}, \frac{\mu - \nu + 2}{2}, 1; \frac{\mu + \nu + 2}{2}, \frac{\mu - \nu + 4}{2}, \frac{3}{2}$$

$$B_2(\mu, \nu) \equiv \frac{\mu + \nu + 2}{2}, \frac{\mu - \nu + 2}{2}, 1; \frac{\mu + \nu + 4}{2}, \frac{\mu - \nu + 4}{2}, \frac{3}{2}$$

$$D_1(\mu) \equiv \frac{1}{2} + \mu, 1; \frac{1}{2} + \mu, \frac{3}{2}$$

$$D_2(\mu) \equiv \frac{1}{2} + \mu, 1; \frac{3}{2} + \mu, \frac{3}{2}$$

$$E_1(\mu, \nu) \equiv \frac{1 + \mu + \nu}{2}, \frac{2 + \mu + \nu}{2}, 1; \frac{2 + \mu + \nu}{2}, \frac{3 + \mu + \nu}{2}, 1 + \nu$$

$$E_2(\mu, \nu) \equiv \frac{1 + \mu + \nu}{2}, \frac{2 + \mu + \nu}{2}, 1; \frac{3 + \mu + \nu}{2}, \frac{4 + \mu + \nu}{2}, 1 + \nu$$

$$F_1(\mu) \equiv \frac{1}{2} + \mu, 1; 1 + \mu, \frac{3}{2} + \mu$$

$$F_2(\mu) \equiv \frac{1}{2} + \mu, 1; 2 + \mu, \frac{3}{2} + \mu$$

REPRESENTATIONS FOR $C_{e_{\mu}}(a, z)$, $C_{s_{\mu}}(a, z)$, $C_{c_{\mu}}(a, z)$

Substituting the Maclaurin series for $\exp(at)$ in Eq. (1) and splitting into even and odd terms we obtain on integrating term by term

$$C_{e_{\mu}}(a, z) = \sum_{n=0}^{\infty} \frac{a^{2n}}{(2n)!} \int_0^z t^{\mu+2n} C_{\nu}(t) dt + \sum_{n=0}^{\infty} \frac{a^{1+2n}}{(1+2n)!} \int_0^z t^{1+\mu+2n} C_{\nu}(t) dt.$$

Then using Eq. (6) and the generating relation Eq. (7) we obtain after a tedious but straightforward computation the principal result of this note

$$\begin{aligned}
 C_{e_{\mu, \nu}}(a, z) = & z^{1+\mu} C_{\nu}(z) \left\{ \frac{1}{\mu - \nu + 1} Q \left[A_1; \frac{a^2 z^2}{4}, \frac{\xi z^2}{4} \right] + \frac{az}{\mu - \nu + 2} Q \left[B_1; \frac{a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right\} \\
 & + \eta z^{2+\mu} C_{\nu-1}(z) \left\{ \frac{1}{(\mu + \nu + 1)(\mu - \nu + 1)} Q \left[A_2; \frac{a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right. \\
 & \left. + \frac{az}{(\mu + \nu + 2)(\mu - \nu + 2)} Q \left[B_2; \frac{a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right\}. \quad (8)
 \end{aligned}$$

Since

$$C_{s_{\mu, \nu}}(a, z) = \frac{1}{2i} \left\{ C_{e_{\mu, \nu}}(ia, z) - C_{e_{\mu, \nu}}(-ia, z) \right\}$$

$$C_{c_{\mu, \nu}}(a, z) = \frac{1}{2} \left\{ C_{e_{\mu, \nu}}(ia, z) + C_{e_{\mu, \nu}}(-ia, z) \right\}$$

we may write

$$\begin{aligned}
 C_{s_{\mu, \nu}}(a, z) = & \frac{az^{2+\mu}}{\mu - \nu + 2} \left\{ C_{\nu}(z) Q \left[B_1; \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right. \\
 & \left. + \frac{\eta z}{\mu + \nu + 2} C_{\nu-1}(z) Q \left[B_2; \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right\} \quad (9)
 \end{aligned}$$

$$\begin{aligned}
 C_{c_{\mu, \nu}}(a, z) = & \frac{z^{1+\mu}}{\mu - \nu + 1} \left\{ C_{\nu}(z) Q \left[A_1; \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right. \\
 & \left. + \frac{\eta z}{\mu + \nu + 1} C_{\nu-1}(z) Q \left[A_2; \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right\}. \quad (10)
 \end{aligned}$$

For $\mu = \nu$, Eqs. (8) to (10) reduce to

$$\begin{aligned}
 C_e(a, z) = & z^{1+\mu} C_{\mu}(z) \left\{ L \left[D_1; \frac{a^2 z^2}{4}, \frac{\xi z^2}{4} \right] + \frac{az}{2} Q \left[B_1(\mu, \mu); \frac{a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right\} \\
 & + \eta z^{2+\mu} C_{\mu-1}(z) \left\{ \frac{1}{1+2\mu} L \left[D_2; \frac{a^2 z^2}{4}, \frac{\xi z^2}{4} \right] + \frac{az}{4(1+\mu)} Q \left[B_2(\mu, \mu); \frac{a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right\} \quad (11)
 \end{aligned}$$

$$\begin{aligned}
 C_s(a, z) = & \frac{1}{2} az^{2+\mu} \left\{ C_{\mu}(z) Q \left[B_1(\mu, \mu); \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right. \\
 & \left. + \frac{\eta z}{2(1+\mu)} C_{\mu-1}(z) Q \left[B_2(\mu, \mu); \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right\} \quad (12)
 \end{aligned}$$

$$C_{\nu}(a, z) = z^{1+\mu} \left\{ C_{\mu}(z) L \left[D_1; \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] + \frac{\eta z}{1+2\mu} C_{\mu-1}(z) L \left[D_2; \frac{-a^2 z^2}{4}, \frac{\xi z^2}{4} \right] \right\}. \quad (13)$$

Defining $J^+ \equiv J$, $J^- \equiv I$, it is interesting to note that we may also write [6]

$$J_{\nu}^{\pm}(a, z) = \frac{z^{1+\mu+\nu} e^{az}}{2^{\nu}(1+\mu+\nu)\Gamma(1+\nu)} \left\{ Q \left[E_1; \frac{\mp z^2}{4}, \frac{a^2 z^2}{4} \right] - \frac{az}{2+\mu+\nu} Q \left[E_2; \frac{\mp z^2}{4}, \frac{a^2 z^2}{4} \right] \right\}$$

$$J_{\nu}^{\pm}(a, z) = \frac{z(z^2/2)^{\mu} e^{az}}{(1+2\mu)\Gamma(1+\mu)} \left\{ L \left[F_1; \frac{\mp z^2}{4}, \frac{a^2 z^2}{4} \right] - \frac{az}{2(1+\mu)} L \left[F_2; \frac{\mp z^2}{4}, \frac{a^2 z^2}{4} \right] \right\}. \quad (14)$$

Here the Bessel functions J_{ν}^{\pm} do not appear.

REDUCTION FORMULAS FOR L AND Q

Many special cases of Eqs. (11) to (14) may be obtained in one form or another, provided we know a reduction formula for either L or Q . We summarize some known relevant reduction formulas [3-6]:

$$L[\alpha, \beta; \gamma, \delta; z, z] = {}_1F_2[\alpha + \beta; \gamma, \delta; z]$$

$$L \left[D_2; \frac{z^2}{4}, \frac{z^2}{4} \right] = \frac{\sinh z}{z}$$

$$L \left[D_1; \frac{z^2}{4}, \frac{z^2}{4} \right] = \frac{2\mu}{1+2\mu} \frac{\sinh z}{z} + \frac{\cosh z}{1+2\mu}$$

$$Q \left[B_2(\mu, \mu); \frac{z^2}{4}, \frac{z^2}{4} \right] = \frac{1+\mu}{1+2\mu} \frac{4}{z^2} \left\{ \cosh z - \left(\frac{2}{z} \right)^{\mu} \Gamma(1+\mu) I_{\mu}(z) \right\}$$

$$Q \left[B_1(\mu, \mu); \frac{z^2}{4}, \frac{z^2}{4} \right] = \frac{2}{1+2\mu} \frac{1}{z} \left\{ 2\mu \frac{\cosh z}{z} + \sinh z - \left(\frac{2}{z} \right)^{\mu} \Gamma(1+\mu) I_{\mu-1}(z) \right\}.$$

Other properties and reduction formulas for L and Q are found in Refs. 3-6.

APPLICATIONS

Of interest in applications are the functions $J_{e_0}(a, z)$, $I_{e_0}(a, z)$, $Y_{e_0}(a, z)$, and $K_{e_0}(a, z)$. $J_{e_0}(a, z)$ and $Y_{e_0}(a, z)$ occur in problems in the theory of diffraction in optical apparatus [1, p. 227]. The function $I_{e_0}(a, z)$ plays an important role in the study of oscillating wings in supersonic flow and arises in the study of resonant absorption in media with finite dimensions [1, p. 195]. $K_{e_0}(a, z)$ occurs when the statistical distribution of the maxima of a random function is applied to the amplitude of a sine wave in order to calculate the distribution of its ordinate. This latter distribution is of

interest in the study of the scattered coherent reflected field from the sea surface [7, 8]. Since the functions $C_{e_0}(a, z)$ are of some importance, by using Eq. (11) and defining

$$L_1(x, y) \equiv L \left[\frac{1}{2}, 1; \frac{1}{2}, \frac{3}{2}; x, y \right]$$

$$L_0(x, y) \equiv L \left[\frac{1}{2}, 1; \frac{3}{2}, \frac{3}{2}; x, y \right]$$

$$Q_1(x, y) \equiv Q \left[1, 1, 1; 1, 2, \frac{3}{2}; x, y \right]$$

$$Q_0(x, y) \equiv Q \left[1, 1, 1; 2, 2, \frac{3}{2}; x, y \right]$$

we obtain

$$\begin{aligned} K_{e_0}(a, z) &= zK_0(z) \left\{ L_1 \left[\frac{a^2 z^2}{4}, \frac{z^2}{4} \right] + \frac{az}{2} Q_1 \left[\frac{a^2 z^2}{4}, \frac{z^2}{4} \right] \right\} \\ &\quad + z^2 K_1(z) \left\{ L_0 \left[\frac{a^2 z^2}{4}, \frac{z^2}{4} \right] + \frac{az}{4} Q_0 \left[\frac{a^2 z^2}{4}, \frac{z^2}{4} \right] \right\} \\ Y_{e_0}(a, z) &= zY_0(z) \left\{ L_1 \left[\frac{a^2 z^2}{4}, \frac{-z^2}{4} \right] + \frac{az}{2} Q_1 \left[\frac{a^2 z^2}{4}, \frac{-z^2}{4} \right] \right\} \\ &\quad + z^2 Y_1(z) \left\{ L_0 \left[\frac{a^2 z^2}{4}, \frac{-z^2}{4} \right] + \frac{az}{4} Q_0 \left[\frac{a^2 z^2}{4}, \frac{-z^2}{4} \right] \right\} \\ J_{e_0}(a, z) &= zJ_0(z) \left\{ L_1 \left[\frac{a^2 z^2}{4}, \frac{-z^2}{4} \right] + \frac{az}{2} Q_1 \left[\frac{a^2 z^2}{4}, \frac{-z^2}{4} \right] \right\} \\ &\quad + z^2 J_1(z) \left\{ L_0 \left[\frac{a^2 z^2}{4}, \frac{-z^2}{4} \right] + \frac{az}{4} Q_0 \left[\frac{a^2 z^2}{4}, \frac{-z^2}{4} \right] \right\} \\ I_{e_0}(a, z) &= zI_0(z) \left\{ L_1 \left[\frac{a^2 z^2}{4}, \frac{z^2}{4} \right] + \frac{az}{2} Q_1 \left[\frac{a^2 z^2}{4}, \frac{z^2}{4} \right] \right\} \\ &\quad - z^2 I_1(z) \left\{ L_0 \left[\frac{a^2 z^2}{4}, \frac{z^2}{4} \right] + \frac{az}{4} Q_0 \left[\frac{a^2 z^2}{4}, \frac{z^2}{4} \right] \right\}. \end{aligned}$$

The equations for $H_{e_0}^{(1)}$ and $H_{e_0}^{(2)}$ are the same as those for Y_{e_0} or J_{e_0} with Y or J replaced by $H^{(1)}$ or $H^{(2)}$. Further, from Eq. (14) we have

$$J_{e_0}(a, z) = ze^{az} \left\{ L \left[1, \frac{1}{2}; \frac{3}{2}, 1; \frac{a^2 z^2}{4}, \frac{-z^2}{4} \right] - \frac{az}{2} L \left[1, \frac{1}{2}; \frac{3}{2}, 2; \frac{a^2 z^2}{4}, \frac{-z^2}{4} \right] \right\}$$

$$I_{e_0}(a, z) = ze^{az} \left\{ L \left[1, \frac{1}{2}; \frac{3}{2}, 1; \frac{a^2 z^2}{4}, \frac{z^2}{4} \right] - \frac{az}{2} L \left[1, \frac{1}{2}; \frac{3}{2}, 2; \frac{a^2 z^2}{4}, \frac{z^2}{4} \right] \right\}.$$

Here we have used the properties of L that

$$L[\alpha, \beta; \gamma, \delta; x, y] = L[\alpha, \beta; \delta, \gamma; x, y] = L[\beta, \alpha; \gamma, \delta; y, x].$$

The latter results for $C_{e_0}(a, z)$ should prove useful in numerical computation of these functions.

SUMMARY

Representations for incomplete Lipschitz-Hankel integrals of cylindrical functions using only the Kampé de Fériet functions in two variables $F_{2;1;0}^{0;2;1}[x, y]$ are given. In addition, known relevant reduction formulas for these functions are provided.

REFERENCES

1. M.M. Agrest and M.S. Maksimov, *Theory of Incomplete Cylindrical Functions and Their Applications* (Springer-Verlag, New York, 1971).
2. Y.L. Luke, *Integrals of Bessel Functions* (McGraw-Hill, New York, 1962).
3. A.R. Miller, "An Incomplete Lipschitz-Hankel Integral of K_0 ," *J. Math. Anal. Appl.* (to appear).
4. A.R. Miller, "A Lipschitz-Hankel Integral of K_0 ," *J. Franklin Institute* (to appear).
5. A.R. Miller, "Incomplete Lipschitz-Hankel Integrals of MacDonald Functions," NRL Report 9112, March 1988.
6. A.R. Miller, "Incomplete Lipschitz-Hankel Integrals of Bessel Functions," *J. Math. Anal. Appl.* (to appear).
7. A.R. Miller and E. Vegh, "A Family of Curves for the Rough Surface Reflection Coefficient," *IEE Proc.-H*, **133**, 483-489 (1986).
8. A.R. Miller and E. Vegh, "Comparison of the Rough Surface Reflection Coefficient with Specularly Scattered Acoustic Data," *J. Acoust. Soc. Amer.*, **82**(5), 1836-1838 (1987).